

BOREL-MOORE HOMOLOGY AND K-THEORY ON THE STEINBERG VARIETY

NAMHEE KWON

1. INTRODUCTION

Let G be a simply connected complex semisimple Lie group with Lie algebra \mathfrak{g} , \mathcal{B} the flag variety of G , \mathcal{N} the nilpotent cone in \mathfrak{g} and $\widetilde{\mathcal{N}}$ the Springer resolution $\{(x, \mathfrak{b}) \in \mathcal{N} \times \mathcal{B} \mid x \in \mathfrak{b}\}$ of \mathcal{N} . We also let

$$Z = \widetilde{\mathcal{N}} \times_{\mathcal{N}} \widetilde{\mathcal{N}} = \{(x, \mathfrak{b}, \mathfrak{b}') \in \mathcal{N} \times \mathcal{B} \times \mathcal{B} \mid x \in \mathfrak{b} \cap \mathfrak{b}'\}$$

be the Steinberg variety. Since its inception, the Steinberg variety has been proved an important object in the representation theory of Weyl groups and affine Hecke algebras. In particular, if we let $H_*(Z, \mathbb{C}) = \bigoplus_{i \geq 0} H_i(Z, \mathbb{C})$ be the complex graded Borel-Moore homology algebra equipped with a convolution product, then the knowledge of the graded algebra structure of $H_*(Z, \mathbb{C})$ was a key ingredient when we obtain all irreducible representations of Weyl groups through the decomposition theorem of Beilinson, Bernstein and Deligne.

In this paper, we study more explicitly the \mathbb{C} -algebra structure of $H_*(Z, \mathbb{C})$. To do so, we will examine the convolution product of $H_*(Z, \mathbb{C})$ through multiplications on another objects such as the Grothendieck group of a graph variety, (co)homology of the flag variety and a certain crossed product algebra. We prove that the convolution product on $H_*(Z, \mathbb{C})$ is compatible with all of these multiplications (see Theorem (3.7)). As an application of our compatibility theorem, we construct a \mathbb{C} -algebra isomorphism between $H_*(Z, \mathbb{C})$ and a certain crossed product algebra (see Theorem (3.9)).

After this paper has been written, the author was informed that Douglass and Röhrle also proved the similar result with Corollary (3.10) (see [DR1, DR2]). However, our approach is different with the one of Douglass and Röhrle.

2. PRELIMINARIES

Borel-Moore homology. Let X be a complex algebraic variety, and let $\widehat{X} = X \cup \{\infty\}$ be the one-point compactification of X . Then the i -th Borel-Moore homology space is defined as $H_i(X) := H_i^{\text{ord}}(\widehat{X}, \infty)$, where H_i^{ord} denotes the i -th singular relative homology over the complex coefficients (see [BM] for details). Throughout the paper, we will assume that all Borel-Moore homology spaces are considered over the complex numbers \mathbb{C} .

2000 *Mathematics Subject Classification.* Primary 20G05; Secondary 18F25, 57T10.

Key words and phrases. Borel-Moore homology, Chern character, K-theory, Steinberg variety.

One of the most important properties of Borel-Moore homology is the existence of a fundamental class $[X]$ for *any* complex algebraic variety X . In particular, let X be a complex algebraic variety of complex dimension n , and let $\{X_1, \dots, X_m\}$ be the n -dimensional irreducible components of X . Then, it is known that the fundamental classes $[X_1], \dots, [X_m]$ form a basis for the vector space $H_{2n}(X)$.

Let M be a smooth variety of complex dimension n . Then, we can consider the following standard \cup -product in cohomology:

$$(2.1) \quad \cup : H^{2n-i}(M) \times H^{2n-j}(M) \longrightarrow H^{4n-i-j}(M).$$

By applying the Poincaré duality to (2.1), we obtain the following bilinear pairing that is called the *intersection pairing*:

$$(2.2) \quad \cap : H_i(M) \times H_j(M) \longrightarrow H_{i+j-2n}(M).$$

Convolution in Borel-Moore homology and K -theory. We first review the construction of the convolution product in Borel-Moore homology due to Ginzburg [CG].

Let M_1, M_2 and M_3 be connected, oriented smooth manifolds with the (i, j) -projections $p_{ij} : M_1 \times M_2 \times M_3 \longrightarrow M_i \times M_j$. We suppose that $Z_{12} \subset M_1 \times M_2$ and $Z_{23} \subset M_2 \times M_3$ are closed subsets.

We also assume that

$$(2.3) \quad p_{13} : \left(p_{12}^{-1}(Z_{12}) \cap p_{23}^{-1}(Z_{23}) \right) \longrightarrow M_1 \times M_3$$

is a proper, and write $Z_{12} \circ Z_{23}$ for the image of p_{13} .

We define a convolution product in Borel-Moore homology as

$$(2.4) \quad * : H_i(Z_{12}) \otimes H_j(Z_{23}) \longrightarrow H_{i+j-d}(Z_{12} \circ Z_{23}), \quad c_{12} \otimes c_{23} \longmapsto c_{12} * c_{23},$$

where $c_{12} * c_{23} = (p_{13})_* \left(p_{12}^* c_{12} \cap p_{23}^* c_{23} \right)$ and $d = \dim_{\mathbb{R}} M_2$.

In particular, if we let $M_1 = M_2 = M_3 = M$ be a smooth manifold of dimension n and $Z_{12} = Z_{23} = M$, then we have the convolution map

$$(2.5) \quad * : H_i(M) \otimes H_j(M) \longrightarrow H_{i+j-2n}(M).$$

In this case, the convolution product in (2.5) reduces to the intersection pairing in (2.2).

A similar convolution construction works for the algebraic K -theory. More explicitly, let X be a smooth algebraic variety and Z_1, Z_2 be closed subsets of X . We also let $K(X)$ be the Grothendieck group of all coherent sheaves on X .

For given $\mathcal{F}_1 \in K(Z_1)$ and $\mathcal{F}_2 \in K(Z_2)$, we write $\mathcal{F}_1 \boxtimes \mathcal{F}_2 \in K(Z_1 \times Z_2)$ for the external tensor product $p_{Z_1}^* \mathcal{F}_1 \otimes_{\mathcal{O}_{Z_1 \times Z_2}} p_{Z_2}^* \mathcal{F}_2$ of \mathcal{F}_1 and \mathcal{F}_2 . Here p_{Z_1} and p_{Z_2} denote the projections of $Z_1 \times Z_2$ to corresponding factors. Next, let $\Delta : X \hookrightarrow X \times X$ be the diagonal embedding. Then we have $\Delta^{-1}(Z_1 \times Z_2) = Z_1 \cap Z_2$. So the restriction map $\Delta : Z_1 \cap Z_2 \longrightarrow Z_1 \times Z_2$ induces the map

$$(2.6) \quad \Delta^* : K(Z_1 \times Z_2) \longrightarrow K(Z_1 \cap Z_2).$$

By combining Δ^* with the external tensor product, we obtain the following tensor product with supports:

$$(2.7) \quad \otimes : K(Z_1) \otimes K(Z_2) \longrightarrow K(Z_1 \cap Z_2), \quad (\mathcal{F}_1, \mathcal{F}_2) \mapsto \Delta^*(\mathcal{F}_1 \boxtimes \mathcal{F}_2).$$

With the tensor product in (2.7), we define the convolution product in the algebraic K -theory as follows.

Let M_1, M_2, M_3 be smooth algebraic varieties. Let $Z_{12} \subset M_1 \times M_2$ and $Z_{23} \subset M_2 \times M_3$ be closed subvarieties satisfying the condition (2.3). Then we define a convolution product

$$(2.8) \quad * : K(Z_{12}) \otimes K(Z_{23}) \longrightarrow K(Z_{12} \circ Z_{23})$$

to be $\mathcal{F}_{12} * \mathcal{F}_{23} := (p_{13})_*((p_{12})^* \mathcal{F}_{12} \otimes (p_{23})^* \mathcal{F}_{23})$.

In particular, the convolution product $* : K(X) \otimes K(X) \longrightarrow K(X)$ coincides with the tensor product $\otimes : K(X) \otimes K(X) \longrightarrow K(X)$ defined in (2.7).

Specializations in Borel-Moore homology and K -theory. Let (X, o) be a smooth manifold with base point o . Given a possibly singular space E , suppose that we have a map $p : E \longrightarrow X$. We also assume that the restriction map $p : P^{-1}(X - o) \longrightarrow X$ is a locally trivial fibration with possibly singular fibers. However, we do not assume that $p : E \longrightarrow X$ is locally trivial around o . According to Fulton and MacPherson [FM], there is a homomorphism

$$\lim_{X \rightarrow o} : H_i(p^{-1}(X - o)) \longrightarrow H_{i - \dim_{\mathbb{R}}(X)}(p^{-1}(o)),$$

which is called a *specialization map*. One of the essential features of the specialization map in Borel-Moore homology is that it commutes with the convolution product of Borel-Moore homology. Similarly, we can define the specialization homomorphism in K -theory. In more detail, let X be a variety and C a smooth algebraic curve with a base point o . Further, let $p : X \longrightarrow C$ be a morphism. Then there exists a specialization homomorphism $\lim_{C \rightarrow o} : K(p^{-1}(X - o)) \longrightarrow K(p^{-1}(o))$. As the case of the specialization map in Borel-Moore homology, the specialization map in K -theory also commutes with the convolution product in K -theory.

The Chern character. Let X be a smooth quasi-projective variety. Then there is a ring homomorphism $ch^* : K(X) \longrightarrow H^*(X)$ called the *cohomological Chern character map* from the Grothendieck ring $K(X)$ to the cohomology ring $H^*(X)$. If we apply the Poincaré duality between the cohomology ring $H^*(X)$ and the graded Borel-Moore homology $H_*(X)$, then we obtain the *homological Chern character map* $ch_* : K(X) \longrightarrow H_*(X)$. Since the cohomological Chern character map $ch^* : K(X) \longrightarrow H^*(X)$ is a ring homomorphism, we see that the homological Chern character map $ch_* : K(X) \longrightarrow H_*(X)$ becomes a ring homomorphism for a smooth quasi-projective variety X (see Corollary (3.4)). The constructions of the Chern character maps over a smooth quasi-projective variety can be extended to those over any closed subvariety of a smooth quasi-projective variety. But, for a non-smooth variety such as the Steinberg variety, we can not guarantee any more that the homological Chern character map is a ring homomorphism.

Throughout the paper, the following notations will be in force:

- $K_{\mathbb{C}}(X)$ denotes the tensor product $\mathbb{C} \otimes_{\mathbb{Z}} K(X)$ for any algebraic variety X .
- \mathcal{B} is the flag variety of a complex semisimple Lie algebra \mathfrak{g} .
- \mathcal{N} is the nilpotent cone in \mathfrak{g} .
- Z denotes the Steinberg variety $\{(x, b_1, b_2) \in \mathcal{N} \times \mathcal{B} \times \mathcal{B} \mid x \in b_1 \cap b_2\}$.

The following theorem shows why we are interested in the homological Chern character map.

Theorem 2.1. *For a variety $X = \mathcal{B}$, $T^*(\mathcal{B})$ or Z , the homological Chern character map yields an isomorphism $K_{\mathbb{C}}(X) \rightarrow H_*(X)$ of abelian groups.*

Proof. See [CG, Theorem 6.2.4]. □

We end up this section with the following theorem.

Theorem 2.2. (1) *The homological Chern character map commutes with the specialization homomorphism in K -theory, and Borel-Moore homology respectively*

(2) *Let X be a smooth variety. Then the following diagram commutes:*

$$(2.9) \quad \begin{array}{ccc} K_{\mathbb{C}}(X) \otimes K_{\mathbb{C}}(X) & \xrightarrow{*= \otimes} & K_{\mathbb{C}}(X) \\ \text{ch}_* \downarrow & & \downarrow \text{ch}_* \\ H_*(X) \otimes H_*(X) & \xrightarrow{*= \cap} & H_*(X). \end{array}$$

Proof. See [BFM]. In fact, in [BFM] we can find a proof of a more generalized fact of (2). Anyway the statement (2) is enough for our purpose, and in this case we will give a simple proof in Corollary (3.4). □

3. \mathbb{C} -ALGEBRA STRUCTURE OF $H_*(Z)$

In the remaining part of this paper, we fix a complex semisimple simply connected Lie group G with the Lie algebra \mathfrak{g} . We also fix a maximal torus T of G .

Universal resolution of \mathfrak{g} . Let $\widetilde{\mathfrak{g}}$ be the universal resolution of \mathfrak{g} (i.e., $\widetilde{\mathfrak{g}} = \{(x, b) \in \mathfrak{g} \times \mathcal{B} \mid x \in b\}$), and let μ be the first projection $\mu : \widetilde{\mathfrak{g}} \rightarrow \mathfrak{g}$. We denote by \widetilde{N} the set $\mu^{-1}(N)$. Since all quotient spaces $b/[b, b]$ are isomorphic for all $b \in \mathcal{B}$, we denote by \mathfrak{S} the the resulting space. We now define a map $\nu : \widetilde{\mathfrak{g}} \rightarrow \mathfrak{S}$ as the quotient map

$$(x, b) \mapsto x + [b, b] \in b/[b, b] \simeq \mathfrak{S}.$$

We write \mathfrak{g}^{sr} for the set of semisimple regular elements in \mathfrak{g} . Here a regular element $x \in \mathfrak{g}$ means that $\dim Z_{\mathfrak{g}}(x) = \text{rank } \mathfrak{g}$. We also denote by $\widetilde{\mathfrak{g}}^{sr}$ the inverse image $\mu^{-1}(\mathfrak{g}^{sr})$. Then, there is a natural action of the Weyl group $W = N(T)/T$ on $\widetilde{\mathfrak{g}}^{sr}$ which preserves fibers $\mu^{-1}(x)$.

Borel isomorphism. Let $W = N_G(T)/T$ be the Weyl group with respect to the maximal torus T . Then the Weyl group W acts on T . This action yields the algebra $\mathbb{C}[T]^W$ of W -invariants, where $\mathbb{C}[T]$ is the algebra of regular functions on T . There is a natural evaluation homomorphism $\mathbb{C}[T]^W \rightarrow \mathbb{C}$ given by evaluation at $1 \in T$.

T . We denote by \mathcal{I} the ideal in $\mathbb{C}[T]$ generated by the kernel of this evaluation homomorphism. It is known from the Borel isomorphism that an algebra $\mathbb{C}[T]/\mathcal{I}$ is isomorphic to the cohomology ring $H^*(\mathcal{B})$ of the flag variety \mathcal{B} as a \mathbb{C} -algebra. We now let $X(T)$ be the character group of T . Then the group algebra $\mathbb{C}[X(T)]$ is identified with $\mathbb{C}[T]$. Through this identification, we will write e^λ for the element of $\mathbb{C}[T]$ corresponding to a weight $\lambda \in X(T)$.

Let $\mathbb{C}[T]/\mathcal{I} \# \mathbb{C}[W]$ be the crossed product of the \mathbb{C} -algebra $\mathbb{C}[T]/\mathcal{I}$ and the group algebra $\mathbb{C}[W]$. Then, this coincides with the tensor product $\mathbb{C}[T]/\mathcal{I} \otimes_{\mathbb{C}} \mathbb{C}[W]$ as a vector space, and it is equipped with multiplication

$$(e^{\lambda_1} \otimes w_1) \cdot (e^{\lambda_2} \otimes w_2) = e^{\lambda_1} (w_1 e^{\lambda_2}) \otimes w_1 w_2,$$

where $w_1 e^{\lambda_2}$ denotes the action of W on $\mathbb{C}[T]$.

Let us now consider the direct sum $\bigoplus_{w \in W} (\mathbb{C}[T]/\mathcal{I}) w$, and define multiplication on it as follows:

$$(e^{\lambda_1} \cdot w_1) (e^{\lambda_2} \cdot w_2) = e^{\lambda_1} (w_1 e^{\lambda_2}) \cdot w_1 w_2.$$

Then it is clear that

$$(3.1) \quad \bigoplus_{w \in W} (\mathbb{C}[T]/\mathcal{I}) w \simeq \mathbb{C}[T]/\mathcal{I} \# \mathbb{C}[W]$$

as \mathbb{C} -algebras.

Graph variety Λ_w^h . From now on, we fix a semisimple regular element h in \mathfrak{g}^{sr} throughout the paper. In [CG], the authors introduced the graph variety $\Lambda_w^h \subset \nu^{-1}(h) \times \nu^{-1}(w \cdot h)$. This is the graph of the action $w : \nu^{-1}(h) \rightarrow \nu^{-1}(w \cdot h)$ obtained by restricting the action $w : \widetilde{\mathfrak{g}}^{sr} \rightarrow \widetilde{\mathfrak{g}}^{sr}$. By the definition of Λ_w^h , we have $\Lambda_w^h = \{(x, \mathfrak{b}, x, Adw(\mathfrak{b})) \in \widetilde{\mathfrak{g}}^{sr} \times \widetilde{\mathfrak{g}}^{sr} \mid x \in \mathfrak{b} \cap Adw(\mathfrak{b}), \nu(x, \mathfrak{b}) = h, \nu(x, Adw(\mathfrak{b})) = Adw(h)\}$.

The following lemma due to Ginzburg.

Lemma 3.1. (1) *The inverse image $\nu^{-1}(h)$ is a smooth variety.*
(2) *The graph variety Λ_w^h is a smooth variety.*

Proof. See [G]. □

Next, let us consider the isomorphisms

$$\nu^{-1}(h) \xrightarrow{w} \nu^{-1}(w \cdot h) \xrightarrow{y} \nu^{-1}(yw \cdot h)$$

for elements $w, y \in W$. Then these isomorphisms yield a set-theoretic composition $\Lambda_w^h \circ \Lambda_y^{w \cdot h}$ defined in (2.3), and we can see that $\Lambda_w^h \circ \Lambda_y^{w \cdot h} = \Lambda_{yw}^h$. Thus we obtain the convolution product

$$(3.2) \quad * : K(\Lambda_w^h) \otimes K(\Lambda_y^{w \cdot h}) \longrightarrow K(\Lambda_{yw}^h).$$

Lemma 3.2. *The map $p : \Lambda_w^h \rightarrow \mathcal{B}$ given by $(x, \mathfrak{b}, x, Adw(\mathfrak{b})) \mapsto \mathfrak{b}$ is a vector bundle over \mathcal{B} with fiber $h + \mathfrak{n}$, and p is a homotopy equivalence.*

Proof. We first notice that Λ_w^h is isomorphic to $\nu^{-1}(h)$, and $\nu^{-1}(h)$ is isomorphic to $G \times_B (h + \mathfrak{n})$ as G -equivariant vector bundles over \mathcal{B} . Moreover, \mathfrak{n} is contractible because it is nilpotent. The lemma now follows. □

The action of the Weyl group W on $H_*(\mathcal{B})$ and $K_{\mathbb{C}}(\mathcal{B})$. Let B be a Borel subgroup containing a maximal torus T , and let W be the Weyl group $W = N_G(T)/T$. We also let $p : G/T \rightarrow G/B = \mathcal{B}$ be the natural projection. Then p is a homotopy equivalence and induces an isomorphism $p^* : H^i(\mathcal{B}) \rightarrow H^i(G/T)$. We notice that the right action of W on G/T given by $w : gT \rightarrow gwT$ yields a W -action on $H^i(G/T)$, and this W -action on $H^i(G/T)$ can be transferred to $H^i(\mathcal{B})$ through the isomorphism p^* . Thus we obtain a W -module structure on $H^i(\mathcal{B})$ which is called the *classical action of W on $H^i(\mathcal{B})$* . We refer [S] for details. Further, if we apply the Poincaré duality $H_*(\mathcal{B}) \simeq H^*(\mathcal{B})$, then the graded Borel-Moore homology $H_*(\mathcal{B})$ also acquires a W -module structure by means of the classical action of W on $H^*(\mathcal{B})$.

Similarly, the homotopy equivalence $p : G/T \rightarrow G/B = \mathcal{B}$ yields the the natural W -module structure on $K_{\mathbb{C}}(\mathcal{B})$ through the right W -action on G/T .

According to [CG, Proposition 6.4.19], there is a W -module isomorphism

$$(3.3) \quad \phi : \mathbb{C}[T]/I \rightarrow K_{\mathbb{C}}(\mathcal{B}).$$

If we combine ϕ with the cohomological Chern character isomorphism $ch^* : K_{\mathbb{C}}(\mathcal{B}) \rightarrow H^*(\mathcal{B})$, then we obtain a W -equivariant algebra isomorphism

$$(3.4) \quad \psi : \mathbb{C}[T]/I \rightarrow H^*(\mathcal{B})$$

which is nothing but the Borel isomorphism introduced in Section 3 (see [CG, Section 6.4] for details).

In the following proposition, we show that for a given smooth quasi-projective variety X the graded Borel-Moore homology $H_*(X)$ is isomorphic to the cohomology ring $H^*(X)$ as \mathbb{C} -algebras.

Proposition 3.3. *Let $H_*(X)$ and $H^*(X)$ be equipped with \mathbb{C} -algebra structures via the convolution product and the cup product respectively. Then the Poincaré duality $H_*(X) \simeq H^*(X)$ yields a \mathbb{C} -algebra isomorphism.*

Proof. We first recall from Section 2 that the convolution $*$: $H_*(X) \otimes H_*(X) \rightarrow H_*(X)$ reduces to the intersection pairing $\cap : H_*(X) \otimes H_*(X) \rightarrow H_*(X)$. However, by definition the intersection pairing $\cap : H_i(X) \otimes H_j(X) \rightarrow H_{i+j-\dim_{\mathbb{R}}X}(X)$ is induced from the standard \cup -product $\cup : H^{\dim_{\mathbb{R}}X-i}(X) \otimes H^{\dim_{\mathbb{R}}X-j}(X) \rightarrow H^{2\dim_{\mathbb{R}}X-i-j}(X)$ through the Poincaré duality. The result now follows. \square

Corollary 3.4. *Let X be a smooth quasi-projective variety. Then the homological Chern character map $ch_* : K_{\mathbb{C}}(X) \rightarrow H_*(X)$ is a \mathbb{C} -algebra homomorphism.*

Proof. We remind that the cohomological Chern character map $ch^* : K_{\mathbb{C}}(X) \rightarrow H^*(X)$ is a ring homomorphism. Then, the result now follows from Proposition (3.3) and the construction of ch_* . \square

Corollary 3.5. *Let $\rho : H^*(\mathcal{B}) \rightarrow H_*(\mathcal{B})$ be the Poincaré duality isomorphism for the flag variety \mathcal{B} . Then the composition map $\rho \circ \psi : \mathbb{C}[T]/I \rightarrow H_*(\mathcal{B})$ is a W -equivariant algebra isomorphism, where ψ is an isomorphism in (3.4).*

Proof. We notice that $\rho : H^*(\mathcal{B}) \rightarrow H_*(\mathcal{B})$ is a W -module isomorphism by the construction of the W -module structure on $H_*(\mathcal{B})$. The result is now immediate from Proposition (3.3) and the W -equivariant algebra isomorphism ψ . \square

Beside the classical action of W on $H_*(\mathcal{B})$, we have another Weyl group action on $H_*(\mathcal{B})$ via the convolution product. In more detail, we first recall from the work of Ginzburg that the group algebra $\mathbb{C}[W]$ is isomorphic to $H_{dim_{\mathbb{R}}Z}(Z)$ as \mathbb{C} -algebras when $H_{dim_{\mathbb{R}}Z}(Z)$ is equipped with the convolution product. We also have the convolution product

$$(3.5) \quad * : H_{dim_{\mathbb{R}}Z}(Z) \otimes H_*(\mathcal{B}) \rightarrow H_*(\mathcal{B})$$

under circumstance $M_1 = M_2 = \tilde{\mathcal{N}}$, $M_3 = \{\text{point}\}$, $Z = \tilde{\mathcal{N}} \times_{\mathcal{N}} \tilde{\mathcal{N}} \subset M_1 \times M_2$ and $\mathcal{B} \simeq \mu^{-1}(0) \subset M_2 \times M_3$, where $\mu : \tilde{\mathfrak{g}} \rightarrow \mathfrak{g}$ is the first projection map.

Then through the identification $\mathbb{C}[W] \simeq H_{dim_{\mathbb{R}}Z}(Z)$ we can define a W -action on $H_*(\mathcal{B})$ via the convolution product in (3.5). Thus we have two W -actions on $H_*(\mathcal{B})$. One is obtained from the classical W -action on $H_*(\mathcal{B})$ and the other comes from the convolution action in (3.5). However, it is known that these two W -actions on $H_*(\mathcal{B})$ are coincide.

Lemma 3.6. *The W -action on $H_*(\mathcal{B})$ arising from the convolution product in (3.5) is the same as the one which is obtained from the classical W -action on $H_*(\mathcal{B})$.*

Proof. See [CG, Claim 3.6.17]. \square

For given $w_1, w_2 \in W$, let

$$\Phi : (\mathbb{C}[T]/\mathcal{I})_{w_1} \otimes (\mathbb{C}[T]/\mathcal{I})_{w_2} \rightarrow K_{\mathbb{C}}(\mathcal{B}) \otimes K_{\mathbb{C}}(\mathcal{B})$$

be a map defined by $\Phi(e^{\lambda}w_1 \otimes e^{\mu}w_2) = \phi(e^{\lambda}) \otimes_{w_1} \phi(e^{\mu})$, where ϕ is the W -module isomorphism in (3.3). Then Φ becomes a vector space isomorphism. We also let $\Psi : (\mathbb{C}[T]/\mathcal{I})_w \rightarrow K_{\mathbb{C}}(\mathcal{B})$ be a W -module isomorphism given by $\Psi(e^{\lambda}w) = \phi(e^{\lambda})$, and write $\rho : H^*(\mathcal{B}) \rightarrow H_*(\mathcal{B})$ for the Poincaré duality isomorphism.

Now, we present the main result of this paper.

Theorem 3.7. *The following diagram commutes for any w_1, w_2 in W :*

$$\begin{array}{ccc}
(\mathbb{C}[T]/I) w_1 \otimes (\mathbb{C}[T]/I) w_2 & \xrightarrow{\text{multiplication}} & (\mathbb{C}[T]/I) w_1 w_2 \\
\Phi \downarrow \wr & & \wr \downarrow \Psi \\
K_{\mathbb{C}}(\Lambda_{w_1}^h) \otimes K_{\mathbb{C}}(\Lambda_{w_2}^{w_1(h)}) & \xrightarrow{*} & K_{\mathbb{C}}(\Lambda_{w_1 w_2}^h) \\
\text{Lemma(3.2)} \downarrow \parallel & & \parallel \downarrow \text{Lemma(3.2)} \\
K_{\mathbb{C}}(\mathcal{B}) \otimes K_{\mathbb{C}}(\mathcal{B}) & \xrightarrow{*= \otimes} & K_{\mathbb{C}}(\mathcal{B}) \\
ch^* \otimes ch^* \downarrow \wr & & \wr \downarrow ch^* \\
H^*(\mathcal{B}) \otimes H^*(\mathcal{B}) & \xrightarrow{\cup} & H^*(\mathcal{B}) \\
\rho \otimes \rho \downarrow \wr & & \wr \downarrow \rho \\
H_*(\mathcal{B}) \otimes H_*(\mathcal{B}) & \xrightarrow{*= \cap} & H_*(\mathcal{B}),
\end{array}$$

where the first row is induced by multiplication in $\bigoplus_{w \in W} (\mathbb{C}[T]/I) w$ and the second row is obtained from (3.2).

Proof. We first recall that the induced K -group homomorphisms from homotopy equivalences commute with the convolution products. From this, we obtain that the second rectangular diagram in Theorem (3.7) commutes.

By definition, the composition map $ch^* \circ \rho : K_{\mathbb{C}}(\mathcal{B}) \rightarrow H_*(\mathcal{B})$ yields the homological Chern character map ch_* .

Then the following diagram commutes due to Theorem (2.2):

$$(3.6) \quad \begin{array}{ccc}
K_{\mathbb{C}}(\mathcal{B}) \otimes K_{\mathbb{C}}(\mathcal{B}) & \xrightarrow{*= \otimes} & K_{\mathbb{C}}(\mathcal{B}) \\
(\rho \otimes \rho) \circ (ch^* \otimes ch^*) = ch_* \otimes ch_* \downarrow \wr & & \wr \downarrow \rho \circ ch^* = ch_* \\
H_*(\mathcal{B}) \otimes H_*(\mathcal{B}) & \xrightarrow{*= \cap} & H_*(\mathcal{B}).
\end{array}$$

Moreover, the definition of the intersection pairing in Section 2 implies that the last rectangular diagram in Theorem (3.7) also commutes. Hence, we obtain the following commutative diagram:

$$(3.7) \quad \begin{array}{ccc}
K_{\mathbb{C}}(\mathcal{B}) \otimes K_{\mathbb{C}}(\mathcal{B}) & \xrightarrow{*= \otimes} & K_{\mathbb{C}}(\mathcal{B}) \\
ch^* \otimes ch^* \downarrow \wr & & \wr \downarrow ch^* \\
H^*(\mathcal{B}) \otimes H^*(\mathcal{B}) & \xrightarrow{\cup} & H^*(\mathcal{B}) \\
\rho \otimes \rho \downarrow \wr & & \wr \downarrow \rho \\
H_*(\mathcal{B}) \otimes H_*(\mathcal{B}) & \xrightarrow{*= \cap} & H_*(\mathcal{B}).
\end{array}$$

Now, we prove that the following diagram commutes which is the main part of this proof:

$$(3.8) \quad \begin{array}{ccc} (\mathbb{C}[T]/\mathcal{I}) w_1 \otimes (\mathbb{C}[T]/\mathcal{I}) w_2 & \xrightarrow{\text{multiplication}} & (\mathbb{C}[T]/\mathcal{I}) w_1 w_2 \\ (\rho \otimes \rho) \circ (ch^* \otimes ch^*) \circ \Phi \Big\downarrow \wr & & \Big\downarrow \wr \rho \circ ch^* \circ \Psi \\ H_*(\mathcal{B}) \otimes H_*(\mathcal{B}) & \xrightarrow{*= \cap} & H_*(\mathcal{B}). \end{array}$$

Let $e^\lambda w_1 \in (\mathbb{C}[T]/\mathcal{I}) w_1$ and $e^\mu w_2 \in (\mathbb{C}[T]/\mathcal{I}) w_2$. Then we have

$$((\rho \otimes \rho) \circ (ch^* \otimes ch^*) \circ \Phi)(e^\lambda w_1 \otimes e^\mu w_2) = (\rho \circ ch^* \circ \phi)(e^\lambda) \otimes w_1 \cdot (\rho \circ ch^* \circ \phi)(e^\mu)$$

because ρ , ch^* and ϕ are W -module isomorphism (see Corollary (3.5) and [CG, Section 6.4]). Notice by Lemma (3.6) that we obtain

$$w_1 \cdot (\rho \circ ch^* \circ \phi)(e^\mu) = [\Lambda_{w_1}] * (\rho \circ ch^* \circ \phi)(e^\mu)$$

if we denote by $[\Lambda_{w_1}]$ the fundamental class corresponding to w_1 through the identification $\mathbb{C}[W] \simeq H_{\dim_{\mathbb{R}} Z}(Z)$. Hence $(\rho \circ ch^* \circ \phi)(e^\lambda) \otimes w_1 \cdot (\rho \circ ch^* \circ \phi)(e^\mu)$ goes to $(\rho \circ ch^* \circ \phi)(e^\lambda) * [\Lambda_{w_1}] * (\rho \circ ch^* \circ \phi)(e^\mu)$ under the convolution product $* : H_*(\mathcal{B}) \otimes H_*(\mathcal{B}) \rightarrow H_*(\mathcal{B})$.

On the other hand, $e^\lambda w_1 \otimes e^\mu w_2$ goes to $e^\lambda (w_1 \cdot e^\mu) w_1 w_2$ under the multiplication map. It now suffices to see the image $(\rho \circ ch^* \circ \phi)(e^\lambda (w_1 \cdot e^\mu))$. We recall from Equation (3.4) that $\psi = ch^* \circ \phi : \mathbb{C}[T]/\mathcal{I} \rightarrow H^*(\mathcal{B})$ and ψ is a W -equivariant algebra isomorphism. So we have

$$\begin{aligned} (\rho \circ ch^* \circ \phi)(e^\lambda (w_1 \cdot e^\mu)) &= \rho(\psi(e^\lambda) \cup w_1 \cdot \psi(e^\mu)) \\ &= (\rho \circ \psi)(e^\lambda) * w_1 \cdot (\rho \circ \psi)(e^\mu) \\ &= (\rho \circ ch^* \circ \phi)(e^\lambda) * [\Lambda_{w_1}] * (\rho \circ ch^* \circ \phi)(e^\mu). \end{aligned}$$

Hence, the diagram in (3.8) commutes.

Since all vertical maps in the diagrams (3.6), (3.7) and (3.8) are isomorphisms, the theorem is now immediate if we combine the commutative diagrams in (3.6), (3.7) and (3.8). \square

For given $w \in W$ and $h \in \mathfrak{g}^{sr}$, let us take the disjoint union set $Z \amalg (\amalg_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h})$. We now consider a map $\pi : Z \amalg (\amalg_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h}) \rightarrow \mathbb{C} \cdot h = \{\lambda h \mid \lambda \in \mathbb{C}\}$ defined by $\pi(Z) = 0$ and $\pi(\Lambda_w^{\lambda h}) = \lambda h$ for each $\lambda \in \mathbb{C}^\times$. Then, the map π induces a locally trivial fibration $\pi : \amalg_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h} \rightarrow \mathbb{C}^\times \cdot h$. Thus we have a specialization homomorphism

$$(3.9) \quad \lim_{\mathbb{C}h \rightarrow 0} : H_* \left(\amalg_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h} \right) \rightarrow H_*(Z).$$

On the other hand, from the homomorphism $\rho \circ ch^* \circ \Psi : (\mathbb{C}[T]/\mathcal{I}) w \rightarrow H_*(\mathcal{B})$ in Theorem (3.7), we obtain a homomorphism

$$(3.10) \quad \theta_w^h : (\mathbb{C}[T]/\mathcal{I}) w \rightarrow H_* \left(\amalg_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h} \right).$$

If we combine the homomorphisms in (3.9) and (3.10), we have a homomorphism

$$(3.11) \quad \theta_w : (\mathbb{C}[T]/\mathcal{I})_w \xrightarrow{\theta_w^h} H_* \left(\coprod_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h} \right) \xrightarrow{\lim_{\mathbb{C}h \rightarrow 0}} H_*(Z),$$

which yields a homomorphism

$$(3.12) \quad \Theta : \bigoplus_{w \in W} (\mathbb{C}[T]/\mathcal{I})_w \longrightarrow H_*(Z)$$

by assembling the homomorphisms θ_w in (3.11). It is known that θ_w does not depend on the choice of h (see [CG, Lemma 3.4.11]).

Corollary 3.8. *The homomorphism Θ in (3.12) is a \mathbb{C} -algebra homomorphism if $H_*(Z)$ is equipped with the convolution product.*

Proof. We recall that the specialization map commutes with the convolution product in Borel-Moore homology. Now the result is immediate from Theorem (3.7) \square

Finally, we construct the following isomorphism.

Theorem 3.9. *The algebra homomorphism $\Theta : \bigoplus_{w \in W} (\mathbb{C}[T]/\mathcal{I})_w \longrightarrow H_*(Z)$ is a \mathbb{C} -algebra isomorphism.*

Proof. It suffices to show that Θ is bijective. We first consider the following diagram:

$$(3.13) \quad \begin{array}{ccc} & \bigoplus_{w \in W} (\mathbb{C}[T]/\mathcal{I})_w & \\ \oplus_{w, \lambda} \Psi \swarrow & & \searrow \oplus_w \theta_w^h \\ \bigoplus_{w \in W} K_{\mathbb{C}} \left(\coprod_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h} \right) & \xrightarrow{\oplus_{w, \lambda} ch_*} & \bigoplus_{w \in W} H_* \left(\coprod_{\lambda \in \mathbb{C}^\times} \Lambda_w^{\lambda h} \right) \\ \oplus \lim^1 \downarrow & & \downarrow \oplus \lim^2 \\ K_{\mathbb{C}}(Z) & \xrightarrow{ch_*} & H_*(Z). \end{array}$$

The diagram in (3.13) is a commutative diagram because

- (1) $\theta_w^h = \bigoplus_{\lambda \in \mathbb{C}^\times} (\rho \circ ch^* \circ \Psi) = \left(\bigoplus_{\lambda \in \mathbb{C}^\times} ch_* \right) \circ \left(\bigoplus_{\lambda \in \mathbb{C}^\times} \Psi \right)$ and
- (2) the homological Chern character commutes with the specialization homomorphism in K -theory, and Borel-Moore homology respectively.

We also notice that the composition map $(\oplus_{w, \lambda} \Psi) \circ (\oplus \lim^1)$ in the diagram (3.13) is a non-equivariant version of the map in [CG, Lemma 7.3.11, 7.3.13]. Thus the map $(\oplus_{w, \lambda} \Psi) \circ (\oplus \lim^1)$ is bijective by applying the same argument in [CG, Lemma 7.3.11, 7.3.13]. The result now follows because the homological Chern character map $ch_* : K_{\mathbb{C}}(Z) \longrightarrow H_*(Z)$ is an isomorphism of abelian groups. \square

Corollary 3.10. *We have \mathbb{C} -algebra isomorphism*

$$\mathbb{C}[T]/\mathcal{I} \# \mathbb{C}[W] \simeq H_*(Z).$$

Proof. Immediate from Equation (3.1) and Theorem (3.9). \square

REFERENCES

- [BFM] P. Baum, W. Fulton and R. MacPherson, *Riemann-Roch for singular varieties*, Publ. Math. IHES **45** (1975), 101-145.
- [BM] A. Borel and J. Moore, *Homology theory for locally compact compact spaces*, Michigan Math. J. **7** (1960), 137-159.
- [CG] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser, Boston, 1997.
- [DR1] J.M. Douglass and G. Röhrle, *Homology of the Steinberg variety and Weyl group coinvariants*, Documenta Math. **14** (2009) 339–357.
- [DR2] J.M. Douglass and G. Röhrle, *The Steinberg variety and representations of reductive groups*, J. Algebra, **321** (2009) 3158-3196.
- [FM] W. Fulton and R. MacPherson, *Categorical framework for the study of singular spaces*, Memoirs of the American Mathematical Society, **243** (1981).
- [G] V. Ginzburg, *\mathfrak{g} -Modules, Springer's representations and bivariant Chern classes*, Advances in Math. **61** (1986), 1-48.
- [S] T. Shoji, *Geometry of orbits and Springer correspondence*, Astérisque. **168** (1988), 61-140.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TOLEDO, 2801 W. BANCROFT ST., TOLEDO, OHIO 43606, USA

E-mail address: namhee.kwon@utoledo.edu